



Hepcidin agonists as therapeutic tools

Carla Casu,¹ Elizabeta Nemeth,² and Stefano Rivella¹

¹Division of Hematology, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, PA; and ²Department of Medicine, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA

Hepcidin agonists are a new class of compounds that regulate blood iron levels, limit iron absorption, and could improve the treatment of hemochromatosis, β -thalassemia, polycythemia vera, and other disorders in which disrupted iron homeostasis causes or contributes to disease. Hepcidin agonists also have the potential to prevent severe complications of siderophilic infections in patients with iron overload or chronic liver disease. This review highlights the preclinical studies that support the development of hepcidin agonists for the treatment of these disorders. (*Blood*. 2018;131(16):1790-1794)

Iron metabolism and the importance of its regulation

Iron is essential for the development and growth of nearly all living organisms, from bacteria to humans.¹ It plays a role in many vital cellular and organismal functions, from cell division to oxygen transport.² As iron is relatively scarce in forms that can be used for biological activities, it is not surprising that organisms developed sophisticated mechanisms for iron acquisition, recycling, and efficient tissue distribution. Although essential, iron in excess can promote the formation of highly toxic reactive oxygen species (ROS), which can damage DNA, protein, and lipid membrane, leading to organ dysfunction.¹⁻³ Therefore, humans and other vertebrates have evolved regulatory systems to optimize the absorption and organ distribution of iron. These regulatory mechanisms are also used to limit iron availability to invading microbes during infection, as microbes compete with the host for this essential nutrient. However, several genetic or acquired disorders of iron homeostasis dysregulate iron absorption or distribution, causing organ damage and creating conditions for overwhelming infections with associated morbidity and mortality.

Hepcidin: the key factor in iron metabolism

The master regulator of iron metabolism is the 25-aa peptide hormone hepcidin (hepcidin antimicrobial peptide [HAMP]), mainly produced by the liver in proportion to plasma iron concentration and iron stores.⁴⁻⁷ Hepcidin inhibits the activity of the only known cellular iron exporter, ferroportin-1 (FPN-1), which is expressed on the surfaces of cells that are involved in iron absorption, recycling, and storage.^{8,9} The feedback circuitry between hepcidin and iron levels in the body ensures systemic iron homeostasis. Hepcidin is also increased during infection and inflammation, causing reduced intestinal iron absorption and increased iron retention in macrophages.¹⁰⁻¹² This is essential to protect the organism from infection with siderophilic and potentially other gram-negative bacteria, which can grow rapidly in the presence of excess iron.¹²⁻¹⁴ In contrast, hepcidin is suppressed

during increased erythropoiesis, when more iron is needed to support increased red blood cell (RBC) production.¹⁵⁻¹⁸

When hepcidin production is chronically decreased, iron absorption and release from body stores are excessive and tissue iron overload occurs.^{6,18} Hepcidin deficiency is a common feature of hereditary hemochromatosis (HH) and anemias with ineffective erythropoiesis, including thalassemias and low-risk myelodysplastic disorders.¹⁹⁻²² In HH, mutations in genes such as *HFE*, *TFR2*, *HJV*, or *HAMP*, which encode proteins in iron-regulatory pathways, lead to reduced hepcidin synthesis and to a phenotype characterized by iron overload from increased intestinal iron absorption and excessive release of iron from macrophages.²² In β -thalassemia, hepcidin deficiency results from chronically elevated production of 1 or more erythroid suppressors of hepcidin.²³⁻²⁸ Mutations in the β -globin gene or its regulatory elements cause reduced or absent β -globin synthesis, resulting in the formation of globin tetramers with an excess of α -chains that precipitate in erythroid progenitors causing their premature death.^{29,30} This, together with the shorter lifespan of RBCs, results in profound anemia that leads to increased erythropoietin production and consequent expansion of the number of erythroblasts, not only in the bone marrow but also at extramedullary sites, causing hepatosplenomegaly.^{19,31} Large numbers of erythroblasts are stimulated by high erythropoietin levels to produce excessive amounts of the erythroid factor erythroferrone, which downregulates hepcidin, leading to increased intestinal iron absorption.²⁷

Utilizing hepcidin pathway to treat iron overload disorders

Initially, studies in mouse models of HH (*Hfe*^{-/-}) or β -thalassemia intermedia (*Hbb*^{th1/th1} and *Hbb*^{th3/+}) demonstrated that transgenic overexpression of hepcidin could prevent iron overload and improve erythropoiesis.^{32,33} Similarly, genetic disruption of the hepcidin regulator *Tmprss6* in a β -thalassemia mouse model enhanced signaling by the bone morphogenetic protein (BMP)/small mothers against decapentaplegic (SMAD) pathway and

Table 1. Classification of hepcidin agonists

Hepcidin agonists	Company	Drug	Target	Clinical trials	Reference
Class 1: hepcidin mimetics	University of California, Los Angeles	MHs (PR65, PR73, M009, M012)	Ferroportin	Validated in preclinical studies	See articles cited in the review
	La Jolla Pharmaceutical Company	LJPC-401 (hepcidin formulation)	Ferroportin	Phase 1: no toxicity reported; expected hypoferrremia observed	45
	Protagonist Therapeutics	PTG-300	Ferroportin	Phase 1: no serious adverse events reported; expected hypoferrremia observed	58
Class 2: stimulators of hepcidin production	Ionis Pharmaceuticals	Tmprss6-ASO	Tmprss6	Phase 1 ongoing	59
	Alynham Pharmaceuticals	Tmprss6-siRNA	Tmprss6	Validated in preclinical studies	See articles cited in the review
Class 3: ferroportin inhibitors	Vifor Pharma	VIT-2763	Ferroportin	Phase 1 planned in 2018	60

ASO, antisense oligonucleotide; siRNA, small-interfering RNA.

increased hepcidin levels with consequent prevention of iron overload and improved erythropoiesis.^{34,35} These genetic studies prompted the exploration of other approaches to increasing circulating hepcidin in iron-overloaded patients with hepcidin deficiency. Synthesis of full-length hepcidin is relatively inefficient, and the half-life of hepcidin in circulation is short because of rapid renal clearance.³⁶ To overcome these limitations, several molecules have been designed to mimic hepcidin activity or stimulate endogenous hepcidin production.^{2,37} We developed minihepcidins (MHs), short peptides based on the 7-9 N-terminal amino acid segment of hepcidin. We first showed that this N-terminal segment of hepcidin is sufficient to induce FPN-1 internalization and degradation in vitro.³⁸ We then engineered these peptides to increase their half-life and potency, and demonstrated that administration of MHs to mice mimics the iron-restrictive effect of endogenous hepcidin.

Studies in a severe mouse model of HH (*Hamp*^{-/-}) showed that the use of MHs could prevent or limit liver iron accumulation and reduce iron level in the heart, while increasing iron sequestration in splenic macrophages.³⁹ Subsequent studies in a mouse model of β -thalassemia intermedia showed that the iron-restrictive effect of MHs improved anemia, iron overload, ineffective erythropoiesis, and splenomegaly.⁴⁰ Treated animals also showed significant reduction in hemichrome formation and ROS as well as improved lifespan of circulating RBCs.⁴⁰ However, this study also demonstrated that high doses of MHs could cause anemia from excessive iron restriction, highlighting the need to titrate hepcidin agonists to the desired effect.⁴⁰

Utilizing hepcidin agonists to treat siderophilic infections

MHs were also highly effective in preventing mortality from infections with siderophilic pathogens in mouse models. Patients with iron-overload disorders are known to be susceptible to severe and lethal infections with siderophilic bacteria such as gram-negative *Vibrio vulnificus* and *Yersinia enterocolitica*.^{41,42} Mouse models of hepcidin deficiency (*Hamp*^{-/-}) reproduced the

susceptibility to lethal infection not only with *V vulnificus* and *Y enterocolitica*, but also with another gram-negative pathogen, *Klebsiella pneumoniae*.^{12-14,43} Even wild-type mice after administration of parenteral iron showed greater infection burden with *Y enterocolitica* and *K pneumoniae*.^{12,14} In mouse models, the pathogenicity of these gram-negative bacteria was dependent on extracellular iron concentrations, specifically on the availability of nontransferrin-bound iron (NTBI). NTBI appears in circulation when transferrin saturation exceeds ~70%, and stimulates extremely rapid proliferation of these bacteria.¹⁴ Administration of MHs to mice rapidly lowered the concentration of extracellular iron, and prevented formation of NTBI. As a result, MHs treatment prevented or reduced mortality from *V vulnificus*, *Y enterocolitica*, and *K pneumoniae* in *Hamp*^{-/-} mice, and decreased abscess formation in iron-loaded wild-type mice infected with *Y enterocolitica*.¹²⁻¹⁴ Based on these studies in mouse models, patients with siderophilic infections in the setting of iron-overload disorders and chronic liver disease associated with high transferrin saturation and the presence of NTBI could benefit from treatment with hepcidin agonists.⁴⁴

Multiple other strategies have been developed to mimic hepcidin activity or increase production of endogenous hepcidin. These include synthetic full-length human hepcidin,⁴⁵ another peptide-based hepcidin mimetic,⁴⁶ small molecule FPN-1 inhibitors,⁴⁷ or pharmacologic inhibition of *Tmprss6* by in vivo targeting and degradation of *Tmprss6* messenger RNA,^{48,49} all displaying similar beneficial effects in preclinical models of hepcidin deficiency. *Tmprss6* antisense oligonucleotide or small-interfering RNA approaches were used in mouse models of β -thalassemia intermedia and hereditary hemochromatosis (*Hfe*^{-/-}).^{48,49} Successful silencing of *Tmprss6* led to increased hepcidin and reduced serum and liver iron concentration in both mouse models. In particular, in β -thalassemia models, increased hepcidin concentration levels were associated with improved anemia, ineffective erythropoiesis (with improved RBC maturation), and reduced splenomegaly. Treated mice also showed reduction in hemichrome formation, apoptosis, and ROS as well as improved RBC life span.^{48,49} Additional studies using MH or *Tmprss6* inhibitors in combination with the iron chelator deferoxamine demonstrated a

more powerful effect of the combined therapy on iron overload and ineffective erythropoiesis than the single agents alone.^{40,50,51}

Hepcidin agonists could also be beneficial for the management of β -thalassemia major (TM). In patients affected by TM, the major cause of iron overload is the iron burden of repeated blood transfusions.^{19,52} Although transfusion therapy transiently increases hepcidin levels, hepcidin decreases toward the end of the transfusion cycle when hemoglobin levels decrease.⁵³ Therefore, increased intestinal iron absorption may also contribute a small but potentially important component of iron overload. In addition, low or relatively low levels of hepcidin in this condition could lead to the generation of NTBI and labile plasma iron, which trigger oxidative stress and tissue injury in many organs.⁴⁴ Therefore, in these disorders, the use of hepcidin agonists could help limit or prevent intestinal iron intake when the synthesis of endogenous hepcidin is not sufficient, and prevent formation of labile plasma iron by sequestering iron in macrophages. Furthermore, the restrictive effect of iron deficiency on ineffective erythropoiesis could also decrease the number of erythroid progenitors and splenomegaly. A smaller spleen should, in turn, reduce the rate of destruction of transfused RBC and reduce the transfusion frequency. To test these hypotheses, we are now conducting a series of studies in a new mouse model of TM.

Another disease in which iron restriction might be beneficial is polycythemia vera (PV).⁴⁰ Most of the clinical characteristics of PV are triggered by the excessive number of erythrocytes, leading to an increased risk of pulmonary hypertension and thrombosis.⁵⁴ The principal therapy for PV is phlebotomy to reduce the hemoglobin concentration and reduce the risk of thrombosis.⁵⁴ Because phlebotomy does not suppress the production of RBCs in the bone marrow, the effect of each phlebotomy is transient, until patients become iron deficient. As an alternative, systemic iron restriction induced by administration of MHs in a mouse model of PV limited iron supply to erythropoiesis and thereby reduced hemoglobin concentrations and decreased splenomegaly.⁴⁰ All of these studies also indicate that overadministration of hepcidin agonists can cause suppression of intestinal iron uptake and macrophage iron recycling, with potential exacerbation of anemia, as in β -thalassemia, or suboptimal production of RBCs, as in PV. These observations stress the notion that careful titration of these compounds is needed to fully benefit from their therapeutic properties, while avoiding undesirable side effects.

Nevertheless, based on the positive results from preclinical studies, several hepcidin agonists are being tested in clinical trials (Table 1).

Conclusions and future directions

Hepcidin agonists have been developed to control iron absorption and ameliorate iron overload in multiple iron disorders,

but their therapeutic efficacy alone or in combination with existing therapies remains to be tested in clinical trials.

In patients with hereditary hemochromatosis, hepcidin agonists may become useful in conjunction with phlebotomy during the treatment phase, or as a stand-alone treatment during the maintenance phase. Furthermore, hepcidin agonists may effectively treat severe siderophilic infections affecting patients with iron overload or chronic liver disease.

Based on preclinical studies, hepcidin agonists could improve not only iron overload but also the anemia in patients with β -thalassemia intermedia. Future studies will explore whether hepcidin agonists could also ameliorate splenomegaly and decrease the need for blood transfusion in individuals affected by β -thalassemia major. Because of their distinctive effect on iron metabolism, hepcidin agonists may also complement other novel treatments that increase RBC production, such as agents that target ligands of the transforming growth factor- β family.⁵⁵⁻⁵⁷

Acknowledgments

The authors thank Tomas Ganz for support, helpful suggestions, and revision of the manuscript.

This work was supported by National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases grants R01 DK107309 (E.N.) and R01 DK095112 and R01 DK090554 (S.R.).

Authorship

Contribution: C.C., E.N., and S.R. wrote and revised the manuscript.

Conflict-of-interest disclosure: E.N. is a consultant and shareholder for Intrinsic LifeSciences and Silarus Therapeutics, and is a consultant for La Jolla Pharmaceutical Company, Keryx Biopharmaceuticals, and Protagonist Therapeutics. S.R. is a consultant for Ionis Pharmaceuticals. C.C. declares no competing financial interests.

ORCID profiles: C.C., 0000-0001-8458-1786; E.N., 0000-0002-3477-2397; S.R., 0000-0002-0938-6558.

Correspondence: Stefano Rivella, Division of Hematology, Department of Pediatrics, Children's Hospital of Philadelphia, Abramson Research Center, Room 316B, 3615 Civic Center Blvd, Philadelphia, PA 19104; e-mail: rivellas@email.chop.edu.

Footnote

Submitted 13 November 2017; accepted 27 February 2018. Prepublished online as *Blood* First Edition paper, 9 March 2018; DOI 10.1182/blood-2017-11-737411.

REFERENCES

- Oikonomidou PR, Casu C, Rivella S. New strategies to target iron metabolism for the treatment of beta thalassemia. *Ann N Y Acad Sci*. 2016;1368(1):162-168.
- Arezes J, Nemeth E. Hepcidin and iron disorders: new biology and clinical approaches. *Int J Lab Hematol*. 2015; 37(suppl 1):92-98.
- Papanikolaou G, Pantopoulos K. Iron metabolism and toxicity. *Toxicol Appl Pharmacol*. 2005;202(2):199-211.
- Park CH, Valore EV, Waring AJ, Ganz T. Hepcidin, a urinary antimicrobial peptide synthesized in the liver. *J Biol Chem*. 2001;276(11):7806-7810.
- Rechavi G, Rivella S. Regulation of iron absorption in hemoglobinopathies. *Curr Mol Med*. 2008;8(7):646-662.
- Ganz T, Nemeth E. Hepcidin and iron homeostasis. *Biochim Biophys Acta*. 2012; 1823(9):1434-1443.
- Muckenthaler MU, Rivella S, Hentze MW, Galy B. A red carpet for iron metabolism. *Cell*. 2017;168(3):344-361.
- Nemeth E, Tuttle MS, Powelson J, et al. Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its

- internalization. *Science*. 2004;306(5704):2090-2093.
9. Kim A, Nemeth E. New insights into iron regulation and erythropoiesis. *Curr Opin Hematol*. 2015;22(3):199-205.
 10. Ganz T, Nemeth E. Iron homeostasis in host defence and inflammation. *Nat Rev Immunol*. 2015;15(8):500-510.
 11. Wang CY, Babitt JL. Hepcidin regulation in the anemia of inflammation. *Curr Opin Hematol*. 2016;23(3):189-197.
 12. Michels KR, Zhang Z, Bettina AM, et al. Hepcidin-mediated iron sequestration protects against bacterial dissemination during pneumonia. *JCI Insight*. 2017;2(6):e92002.
 13. Arezes J, Jung G, Gabayan V, et al. Hepcidin-induced hypoferremia is a critical host defense mechanism against the siderophilic bacterium *Vibrio vulnificus*. *Cell Host Microbe*. 2015;17(1):47-57.
 14. Stefanova D, Raychev A, Arezes J, et al. Endogenous hepcidin and its agonist mediate resistance to selected infections by clearing non-transferrin-bound iron. *Blood*. 2017;130(3):245-257.
 15. Nicolas G, Chauvet C, Viatte L, et al. The gene encoding the iron regulatory peptide hepcidin is regulated by anemia, hypoxia, and inflammation. *J Clin Invest*. 2002;110(7):1037-1044.
 16. Nemeth E, Rivera S, Gabayan V, et al. IL-6 mediates hypoferremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest*. 2004;113(9):1271-1276.
 17. Pak M, Lopez MA, Gabayan V, Ganz T, Rivera S. Suppression of hepcidin during anemia requires erythropoietic activity. *Blood*. 2006;108(12):3730-3735.
 18. Oikonomidou PR, Rivella S. What can we learn from ineffective erythropoiesis in thalassemia? *Blood Rev*. 2017;S0268-960X(17)30131-5.
 19. Ginzburg Y, Rivella S. β -thalassemia: a model for elucidating the dynamic regulation of ineffective erythropoiesis and iron metabolism. *Blood*. 2011;118(16):4321-4330.
 20. Camaschella C, Nai A. Ineffective erythropoiesis and regulation of iron status in iron loading anaemias. *Br J Haematol*. 2016;172(4):512-523.
 21. Sebastiani G, Wilkinson N, Pantopoulos K. Pharmacological targeting of the hepcidin/ferroportin axis. *Front Pharmacol*. 2016;7:160.
 22. Powell LW, Seckington RC, Deugnier Y. Haemochromatosis. *Lancet*. 2016;388(10045):706-716.
 23. Gardenghi S, Marongiu MF, Ramos P, et al. Ineffective erythropoiesis in beta-thalassemia is characterized by increased iron absorption mediated by down-regulation of hepcidin and up-regulation of ferroportin. *Blood*. 2007;109(11):5027-5035.
 24. Papanikolaou G, Tzilianos M, Christakis JI, et al. Hepcidin in iron overload disorders. *Blood*. 2005;105(10):4103-4105.
 25. Weizer-Stern O, Adamsky K, Amariglio N, et al. Downregulation of hepcidin and haemojuvelin expression in the hepatocyte cell-line HepG2 induced by thalassaemic sera. *Br J Haematol*. 2006;135(1):129-138.
 26. Weizer-Stern O, Adamsky K, Amariglio N, et al. mRNA expression of iron regulatory genes in beta-thalassemia intermedia and beta-thalassemia major mouse models. *Am J Hematol*. 2006;81(7):479-483.
 27. Kautz L, Jung G, Valore EV, Rivella S, Nemeth E, Ganz T. Identification of erythroferrone as an erythroid regulator of iron metabolism. *Nat Genet*. 2014;46(7):678-684.
 28. Sonnweber T, Nachbauer D, Schroll A, et al. Hypoxia induced downregulation of hepcidin is mediated by platelet derived growth factor BB. *Gut*. 2014;63(12):1951-1959.
 29. Schrier SL. Pathophysiology of thalassemia. *Curr Opin Hematol*. 2002;9(2):123-126.
 30. Rivella S. β -thalassemias: paradigmatic diseases for scientific discoveries and development of innovative therapies. *Haematologica*. 2015;100(4):418-430.
 31. Melchiori L, Gardenghi S, Rivella S. beta-Thalassemia: HiJAKing ineffective erythropoiesis and iron overload. *Adv Hematol*. 2010;2010:938640.
 32. Nicolas G, Viatte L, Lou DQ, et al. Constitutive hepcidin expression prevents iron overload in a mouse model of hemochromatosis. *Nat Genet*. 2003;34(1):97-101.
 33. Gardenghi S, Ramos P, Marongiu MF, et al. Hepcidin as a therapeutic tool to limit iron overload and improve anemia in β -thalassemic mice. *J Clin Invest*. 2010;120(12):4466-4477.
 34. Finberg KE, Whittlesey RL, Andrews NC. Tmprss6 is a genetic modifier of the Hfe-hemochromatosis phenotype in mice. *Blood*. 2011;117(17):4590-4599.
 35. Nai A, Pagani A, Mandelli G, et al. Deletion of Tmprss6 attenuates the phenotype in a mouse model of β -thalassemia. *Blood*. 2012;119(21):5021-5029.
 36. Schmidt PJ, Fleming MD. Modulation of hepcidin as therapy for primary and secondary iron overload disorders: preclinical models and approaches. *Hematol Oncol Clin North Am*. 2014;28(2):387-401.
 37. Crielgaard BJ, Lammers T, Rivella S. Targeting iron metabolism in drug discovery and delivery. *Nat Rev Drug Discov*. 2017;16(6):400-423.
 38. Preza GC, Ruchala P, Pinon R, et al. Minihepcidins are rationally designed small peptides that mimic hepcidin activity in mice and may be useful for the treatment of iron overload. *J Clin Invest*. 2011;121(12):4880-4888.
 39. Ramos E, Ruchala P, Goodnough JB, et al. Minihepcidins prevent iron overload in a hepcidin-deficient mouse model of severe hemochromatosis. *Blood*. 2012;120(18):3829-3836.
 40. Casu C, Oikonomidou PR, Chen H, et al. Minihepcidin peptides as disease modifiers in mice affected by β -thalassemia and polycythemia vera. *Blood*. 2016;128(2):265-276.
 41. Vadillo M, Corbella X, Pac V, Fernandez-Viladrich P, Pujol R. Multiple liver abscesses due to *Yersinia enterocolitica* discloses primary hemochromatosis: three cases reports and review. *Clin Infect Dis*. 1994;18(6):938-941.
 42. Khan FA, Fisher MA, Khakoo RA. Association of hemochromatosis with infectious diseases: expanding spectrum. *Int J Infect Dis*. 2007;11(6):482-487.
 43. Horseman MA, Surani S. A comprehensive review of *Vibrio vulnificus*: an important cause of severe sepsis and skin and soft-tissue infection. *Int J Infect Dis*. 2011;15(3):e157-e166.
 44. Brissot P, Ropert M, Le Lan C, Loréal O. Non-transferrin bound iron: a key role in iron overload and iron toxicity. *Biochim Biophys Acta*. 2012;1820(3):403-410.
 45. La Jolla Pharmaceutical Company. LJPC-401. <http://lajollapharmaceutical.com/product-pipeline/ljpc-401/>. Accessed 14 February 2018.
 46. Bourne G, Zhao L, Bhandari A, et al. Hepcidin mimetic PTG-300 for treatment of ineffective erythropoiesis and iron overload. In: Seventh Congress of the International Biolron Society (IBIS) Biennial World Meeting (Biolron 2017) May 7-11, 2017 [abstract]. *Am J Hematol*. 2017;92(8):E223.
 47. Manolova V, Flace A, Altermatt P, Sundström H, Nyffenegger N, Duerrenberger F. Ferroportin inhibitors improve ineffective erythropoiesis and prevent iron loading in a beta-thalassemia disease model. In: Seventh Congress of the International Biolron Society (IBIS) Biennial World Meeting (Biolron 2017) May 7-11, 2017 [abstract]. *Am J Hematol*. 2017;92(8):E224.
 48. Guo S, Casu C, Gardenghi S, et al. Reducing Tmprss6 ameliorates hemochromatosis and β -thalassemia in mice. *J Clin Invest*. 2013;123(4):1531-1541.
 49. Schmidt PJ, Toudjarska I, Sendamarai AK, et al. An RNAi therapeutic targeting Tmprss6 decreases iron overload in Hfe(-/-) mice and ameliorates anemia and iron overload in murine β -thalassemia intermedia. *Blood*. 2013;121(7):1200-1208.
 50. Schmidt PJ, Racie T, Westerman M, Fitzgerald K, Butler JS, Fleming MD. Combination therapy with a Tmprss6 RNAi-therapeutic and the oral iron chelator deferiprone additively diminishes secondary iron overload in a mouse model of β -thalassemia intermedia. *Am J Hematol*. 2015;90(4):310-313.
 51. Casu C, Aghajan M, Oikonomidou PR, Guo S, Monia BP, Rivella S. Combination of Tmprss6-ASO and the iron chelator deferiprone improves erythropoiesis and reduces iron overload in a mouse model of beta-thalassemia intermedia. *Haematologica*. 2016;101(1):e8-e11.
 52. Origa R, Galanello R, Ganz T, et al. Liver iron concentrations and urinary hepcidin in beta-thalassemia. *Haematologica*. 2007;92(5):583-588.
 53. Pasricha SR, Frazer DM, Bowden DK, Anderson GJ. Transfusion suppresses erythropoiesis and increases hepcidin in adult patients with β -thalassemia major: a longitudinal study. *Blood*. 2013;122(1):124-133.

54. Silver RT, Gjoni S. The hematocrit value in polycythemia vera: caveat utilitor. *Leuk Lymphoma*. 2015;56(5):1540-1541.
55. Breda L, Rivella S. Modulators of erythropoiesis: emerging therapies for hemoglobinopathies and disorders of red cell production. *Hematol Oncol Clin North Am*. 2014;28(2):375-386.
56. Suragani RN, Cawley SM, Li R, et al. Modified activin receptor IIB ligand trap mitigates ineffective erythropoiesis and disease complications in murine β -thalassemia. *Blood*. 2014;123(25):3864-3872.
57. Dussiot M, Maciel TT, Fricot A, et al. An activin receptor IIA ligand trap corrects ineffective erythropoiesis in β -thalassemia. *Nat Med*. 2014;20(4):398-407.
58. Protagonist Therapeutics. Protagonist Therapeutics announces final phase 1 study results with novel hepcidin mimetic, PTG-300. <https://www.pnewswire.com/news-releases/protagonist-therapeutics-announces-final-phase-1-study-results-with-novel-hepcidin-mimetic-ptg-300-300571268.html>. Accessed 14 February 2018.
59. Ionis Pharmaceuticals. Pipeline. <http://www.ionispharma.com/pipeline/>. Accessed 14 February 2018.
60. Vifor Pharma. Vifor Pharma ferroportin inhibitor enters phase-I clinical trial. <http://www.viforpharma.com/en/media/press-releases/201802/2167159>. Accessed 14 February 2018.